



**SURGICAL AND MEDICAL SUPPLIES PTY. LTD.**

ABN 51 007 707 128

16 Kensington Road Rose Park South Australia 5067

PO Box 2 Kent Town SA 5071

Telephone: (08) 8332 1666 Facsimile: (08) 8332 5132

www.surmed.com.au Email: sales@surmed.com.au

**FITTINGS BY APPOINTMENT ONLY – PH 8332 1666**

Date: \_\_\_\_\_

Please fit my patient with a SIGVARIS® graduated compression garment.

Name: \_\_\_\_\_

Street: \_\_\_\_\_

Suburb: \_\_\_\_\_ Postcode: \_\_\_\_\_

**DIAGNOSIS:** \_\_\_\_\_  
\_\_\_\_\_

Compression Class				
Travel Socks	(Y / N)			
Calf	1	2	3	4
Thigh	1	2	3	4
Thigh (with waist attachment)	1	2	3	
Pantyhose	1	2		
Maternity Pantyhose	1	2		
Arm Sleeve	1	2		
Arm Sleeve (with Mitten)	1	2		

Please circle above the compression class needed

(Compression Classes: 1 = 18-21mmHg, 2 = 23-32mmHg, 3 = 34-46mmHg, 4 = min 49mmHg)

Travel socks = 15-18mmHg



Practitioners Name: \_\_\_\_\_

Signature: \_\_\_\_\_

Address: \_\_\_\_\_

Phone: \_\_\_\_\_

# Indications

# Contraindications

INDICATIONS	C.CL RECOMMENDED
Prophylaxis of venous disorders in patients at risk	1
Functional disorders such as heaviness and fatigue in the legs, with existing varicosis (C <sub>0</sub> ) <sup>▲</sup>	1
Edema in the evening, due to local causes (venous) (C <sub>3</sub> ) <sup>▲</sup>	1-2
Varicosis during pregnancy, without edema (C <sub>2</sub> ) <sup>▲</sup>	1-2
Varicosis during pregnancy, with edema (C <sub>3</sub> ) <sup>▲</sup>	2-3
Mild varices with subjective manifestations (small varicose dilatation of the cutaneous veins, reticular varices) (C <sub>1</sub> -C <sub>2</sub> ) <sup>▲</sup>	1
Primary varicosis with edema (C <sub>3</sub> ) <sup>▲</sup>	2
Advanced primary varicosis with edema (stem and branch varices) (C <sub>3</sub> ) <sup>▲</sup>	2-3
Chronic venous insufficiency (CVI)	
Grade I* (C <sub>3</sub> ) <sup>▲</sup> corona phlebectatica paraplantaris	2
Grade II* (C <sub>4</sub> ) <sup>▲</sup> chronic edema eczema erythema hypodermatitis dermatosclerosis dermite ocre atrophie blanche	depending on severity after completion of basic therapy 2 or 3
Grade IIIa* (C <sub>5</sub> ) <sup>▲</sup> healed ulcer cruris (relapse prophylaxis)	3
Post-traumatic edema after completion of basic therapy (C <sub>3</sub> ) <sup>▲</sup>	3 (possibly also 2)
Prophylaxis of thrombosis/embolism	1-2
Thrombophlebitis (superficial thrombosis), possibly with edema	2
Thrombophlebitis (deep-vein thrombosis) after completion of basic therapy	depending on severity 2 or 3
Post-thrombotic edema with or without secondary follow-up therapy	depending on severity 2 or 3
Angiodysplasia	depending on involvement 2, 3 or 4
Before sclerotherapy or before surgery	2
To maintain success of therapy: a) after sclerotherapy; b) after ambulant selective varicose surgery; c) after stripping	2
Reversible lymphedema after complex physical decongestant therapy	3
Irreversible lymphedema with pronounced induration after complex physical decongestant therapy	4
Lipedema, after completion of basic therapy	4

\* according to Widmer & Marshall ▲ according to CEAP-Classification

**In case of overweight, prescribe the next higher compression category.**

**ABSOLUTE CONTRAINDICATIONS** Advanced peripheral arterial occlusive disease, Congestive heart failure, Septic phlebitis (inflammation of the vein caused by microorganisms), Phlegmasia coerulea dolens<sup>1</sup>

**RELATIVE CONTRAINDICATIONS** Oozing dermatoses, Intolerance of compression stocking material, Sensory loss of the extremity, Advanced peripheral neuropathy (e.g. diabetes mellitus), Rheumatoid arthritis (RA)

**No liability accepted in case of non-observance of these contraindications.**