



SURGICAL AND MEDICAL SUPPLIES PTY. LTD.

ABN 51 007 707 128

16 Kensington Road Rose Park South Australia 5067

PO Box 2 Kent Town SA 5071

Telephone: (08) 8332 1666 Facsimile: (08) 8332 5132

www.surmed.com.au Email: sales@surmed.com.au

**PLEASE PH 8332 1666 FOR A FITTING FORM
or download one from our website (search for Sigvaris)**

Date: _____

Please fit my patient with a SIGVARIS® graduated compression garment.

Name: _____

Street: _____

Suburb: _____ Postcode: _____

DIAGNOSIS: _____

	Compression Class			
Travel Socks	(Y / N)			
Calf	1	2	3	4
Thigh	1	2	3	4
Thigh (with waist attachment)	1	2	3	
Pantyhose	1	2		
Maternity Pantyhose	1	2		
Arm Sleeve	1	2		
Arm Sleeve (with Mitten)	1	2		

Please circle above the compression class needed

(Compression Classes: 1 = 18-21mmHg, 2 = 23-32mmHg, 3 = 34-46mmHg, 4 = min 49mmHg)

Travel socks = 15-18mmHg



Practitioners Name: _____

Signature: _____

Address: _____

Phone: _____

Indications

Contraindications

INDICATIONS	C.CL RECOMMENDED
Prophylaxis of venous disorders in patients at risk	1
Functional disorders such as heaviness and fatigue in the legs, with existing varicosis (C ₀) [▲]	1
Edema in the evening, due to local causes (venous) (C ₃) [▲]	1-2
Varicosis during pregnancy, without edema (C ₂) [▲]	1-2
Varicosis during pregnancy, with edema (C ₃) [▲]	2-3
Mild varices with subjective manifestations (small varicose dilatation of the cutaneous veins, reticular varices) (C ₁ -C ₂) [▲]	1
Primary varicosis with edema (C ₃) [▲]	2
Advanced primary varicosis with edema (stem and branch varices) (C ₃) [▲]	2-3
Chronic venous insufficiency (CVI)	
Grade I* (C ₃) [▲] corona phlebectatica paraplanaris	2
Grade II* (C ₄) [▲] chronic edema eczema erythema hypodermatitis dermatosclerosis dermite ocre atrophie blanche	depending on severity after completion of basic therapy 2 or 3
Grade IIIa* (C ₅) [▲] healed ulcer cruris (relapse prophylaxis)	3
Post-traumatic edema after completion of basic therapy (C ₃) [▲]	3 (possibly also 2)
Prophylaxis of thrombosis/embolism	1-2
Thrombophlebitis (superficial thrombosis), possibly with edema	2
Thrombophlebitis (deep-vein thrombosis) after completion of basic therapy	depending on severity 2 or 3
Post-thrombotic edema with or without secondary follow-up therapy	depending on severity 2 or 3
Angiodysplasia	depending on involvement 2, 3 or 4
Before sclerotherapy or before surgery	2
To maintain success of therapy: a) after sclerotherapy; b) after ambulant selective varicose surgery; c) after stripping	2
Reversible lymphedema after complex physical decongestant therapy	3
Irreversible lymphedema with pronounced induration after complex physical decongestant therapy	4
Lipedema, after completion of basic therapy	4

* according to Widmer & Marshall ▲ according to CEAP-Classification

In case of overweight, prescribe the next higher compression category.

ABSOLUTE CONTRAINDICATIONS Advanced peripheral arterial occlusive disease, Congestive heart failure, Septic phlebitis (inflammation of the vein caused by microorganisms), Phlegmasia coerulea dolens¹

RELATIVE CONTRAINDICATIONS Oozing dermatoses, Intolerance of compression stocking material, Sensory loss of the extremity, Advanced peripheral neuropathy (e.g. diabetes mellitus), Rheumatoid arthritis (RA)

No liability accepted in case of non-observance of these contraindications.